


# **Climate Change Brief**

*City and County of Honolulu – Climate Change Commission*

Adopted: June 5, 2018

The Climate Change Commission consists of five members with expertise in climate change in Hawai'i. The Climate Change Commission was created after O'ahu voters approved an amendment to the City Charter in the 2016 general election that also mandated the creation of the Office of Climate Change, Sustainability and Resiliency. The role of the Climate Change Commission is to gather the latest science and information on climate change impacts to Hawai'i and provide advice and recommendations to the mayor, City Council, and executive departments as they look to draft policy and engage in planning for future climate scenarios.

<https://www.resilientoahu.org/about-the-commission/>



**Figure 1.** Global temperature since 1880. In 1970 global surface temperature began a consistent, long-term rise at a rate of 0.3°F (0.17°C) per decade. Individual years, or groups of years, may be warmer or cooler than this trend due to various causes of climate variability. These include cooling effects from light scattering particles in the atmosphere (e.g., sulfur aerosols from volcanic and industrial emissions), the El Niño Southern Oscillation, the Pacific Decadal Oscillation, and others.

Source: Data from NASA Goddard Institute for Space Studies: <https://data.giss.nasa.gov/gistemp/graphs/>; figure from [http://www.columbia.edu/~jeh1/mailings/2018/20180118\\_Temperature2017.pdf](http://www.columbia.edu/~jeh1/mailings/2018/20180118_Temperature2017.pdf)

## KEY POINTS

- 1) Although greenhouse gas emissions have only warmed Earth's surface 1.8°F (1°C), the impacts are already severe and widespread, in part because they magnify existing stress that human communities place on natural systems.
- 2) Climate change effects are documented across every ecosystem on Earth and scientists have concluded that a sixth mass extinction is underway. The ocean is hotter and more acidic with less dissolved O<sub>2</sub>. A rise in extreme weather has been documented including, expanded drought and desertification, record setting intense rainfall, a rise in extreme hurricanes, food and water shortages, increased political turmoil, and tens of millions of internally and externally displaced refugees.
- 4) In Hawai'i, documented effects include rising air temperature that is warming four times faster than half a century ago. Decreasing wind speeds and rainfall in dry and wet seasons. A rise in extreme rain and drought. Decreased stream flow indicating declining groundwater levels. Mosquito-borne diseases are migrating to previously safe upland forests. Severe ocean changes leading to increased coral bleaching. Long-term sea level rise driving chronic beach erosion and loss.
- 5) In Hawai'i, these trends are all expected to accelerate as the air continues to warm. Future projections include more frequent tropical cyclones, and extreme El Niño events, 50% of native forest bird species will lose over half their range, and as early as 2030, almost all reefs are expected to experience annual bleaching leading to their extinction.
- 6) With 3.2 ft (0.98 m) of sea level rise this century, 25,800 acres in Hawai'i will experience chronic flooding, erosion, and high wave impacts. One third of this land is designated for urban use. Impacts include 38 mi (61 km) of major roads, and more than \$19 billion in assets.
- 5) The negative impacts of climate change fall disproportionately on disadvantaged groups.
- 6) This information reinforces the urgent need for a decarbonized world economy. Nothing less than a sweeping transformation in our energy sources, land-use practices, and food and water systems will stop continued warming.
- 7) However, economists project a 15% increase in carbon dioxide emissions over the next 20 years, setting us on a pathway to a world that is more than 5.4°F (3°C) warmer. This commits Hawai'i to a challenging future that requires strong efforts to increase community resilience and sustainability.

## PURPOSE

Pursuant to the Revised Charter of Honolulu (“RCH”) Section 6-107(h), the City and County of Honolulu (“City”) Climate Change Commission is charged with gathering the latest science and information on climate change impacts to Hawai‘i and providing advice and recommendations to the mayor, City Council, and executive departments as they look to draft policy and engage in planning for future climate scenarios and reducing Honolulu’s contribution to global greenhouse gas emissions.

To establish the factual basis and broad impact of climate change, the City Climate Change Commission adopts this CLIMATE CHANGE BRIEF 2018. This document describes the local, regional, and global impacts of climate change as documented by the peer-reviewed scientific literature and credible empirical data sources. It provides a benchmark for the commission, attesting to our concern, underpinning our decisions and recommendations, and serving to inform those we serve.

The information in this report reinforces the need for an urgent and sweeping transformation in our energy sources, food systems, and land-use practices to achieve a decarbonized world economy. Mitigation of future climate change must be achieved to avoid the very worst aspects of global warming. In the words of Dr. Jim Hansen, former chief scientist at the NASA Goddard Institute of Space Science, “There is a possibility, a real danger, that we will hand young people and future generations a climate system that is practically out of their control... we have a global emergency. Fossil fuel CO<sub>2</sub> emissions should be reduced as rapidly as practical.”<sup>1</sup>

Because many changes in global biogeochemical systems have been irreversibly set into motion, and these threaten the health and welfare of human populations, it is important that the City and County of Honolulu take bold steps to reduce greenhouse gas emissions and build sustainability and resilience in the face of a rapidly changing climate.

## INTRODUCTION

Excess heat, trapped by the anthropogenic greenhouse gases carbon dioxide, methane, nitrous oxide, and others in the atmosphere, is causing dramatic changes in ecosystems, the ocean, weather patterns, and other climate-dependent aspects of Earth’s surface. Hawai‘i, and other Pacific islands are impacted, and these impacts are growing.<sup>2</sup>

The negative impacts of climate change fall disproportionately on disadvantaged groups in a type of “vicious cycle”.<sup>3</sup> Initial inequity or vulnerabilities can be exacerbated by climate change; for example, low income people are less likely to have air conditioning and can be much more susceptible to the effects of a heat wave. This in turn lowers the ability of already disadvantaged groups to cope and recover. It is important to recognize and resolve the impacts of climate change on vulnerable populations as the City pivots to meet the challenges of climate change.

Unrelenting impacts to Earth’s ecosystems<sup>4</sup> and natural resources have led researchers to conclude that our planet is perched on the edge of a tipping point<sup>5</sup>, a planetary-scale critical transition resulting from human impacts.<sup>6</sup> These changes include the following.

### CARBON DIOXIDE

- Carbon dioxide levels in the air have passed 410 ppm compared to a natural level of 280 ppm<sup>7</sup> – an increase of over 45%. This is the highest level in millions of years.<sup>8</sup>
- Today, release of planet-warming carbon dioxide is ten times faster than the most rapid event in the past 66 million years, when an asteroid impact killed the dinosaurs.<sup>9</sup>

### TEMPERATURE

- Global temperature has risen approximately 1.8°F (1°C) from the late 19<sup>th</sup> Century.<sup>10</sup>
- The likely global temperature increase this century is a median 5.76°F (3.2°C). There is only a 5% chance that it will be less than 3.6°F (2°C), and a 1% chance that it will

be less than 2.7°F (1.5°C).<sup>11</sup>

- The last time it was this warm, 125,000 years ago, global sea level was 20 ft (6.6 m) higher.<sup>12 13 14</sup>
- Atmospheric humidity is rising.<sup>15</sup>
- The global water cycle has accelerated.<sup>16</sup>
- Air temperature over the oceans is rising.<sup>17</sup>

### HAWAI‘I – LOCAL AND REGIONAL IMPACTS

#### *Air Temperature*

- In Hawai‘i, the rate of warming air temperature has increased in recent decades. Currently, the air is warming at 0.3°F (0.17°C) per decade, four times faster than half a century ago.<sup>18</sup>
- Statewide, average air temperature has risen by 0.76°F (0.42°C) over the past 100 years, and 2015 and 2016 were the warmest years on record.<sup>19</sup>

- Warming air temperatures lead to heat waves, expanded pathogen ranges and invasive species, thermal stress for native flora and fauna, increased electricity demand, increased wildfire, potential threats to human health, and increased evaporation which both reduces water supply and increases demand. Rapid warming at highest elevations impedes precipitation, the source of Hawai'i's freshwater.<sup>20</sup>
- During the strong El Niño of 2015, Honolulu set or tied 11 days of record heat.<sup>21</sup> This compelled the local energy utility to issue emergency public service announcements to curtail escalating air conditioning use that stressed the electrical grid.<sup>22</sup>
- Some model projections for the late 21<sup>st</sup> century indicate that surface air temperature over land will increase 1.8° to 7.2°F (2° to 4°C) with the greatest warming at the highest elevations and on leeward sides of the major islands.<sup>23</sup>
- Under continued strong greenhouse gas emissions, high elevations above 9,800 ft (3000 m) reach up to 7.2° to 9°F (4 to 5°C) warmer temperatures by the late 21<sup>st</sup> Century.<sup>24</sup>

### **Wind and Precipitation**

- The frequency of gale-force winds is increasing in the western and south Pacific but decreasing in the central Pacific.<sup>25</sup>
- Average daily wind speeds are slowly declining in Honolulu and Hilo, while remaining steady across western and south Pacific sites.<sup>26</sup>
- Studies indicate there will be future changes to winds and waves due to climate change, which affects ecosystems, infrastructure, freshwater availability, and commerce.<sup>27</sup>
- Hawai'i has seen an overall decline in rainfall over the past 30 years, with widely varying precipitation patterns on each island. The period since 2008 has been particularly dry.<sup>28</sup>
- Declining rainfall has occurred in both the wet and dry seasons and has affected all the major islands. On O'ahu, the largest declines have occurred in the northern Ko'olau mountains.<sup>29</sup>
- Heavy rainfall events and droughts have become more common, increasing runoff, erosion, flooding, and water shortages.<sup>30</sup>
- Consecutive wet days and consecutive dry days are both increasing in Hawai'i.<sup>31</sup>
- There is disagreement regarding precipitation at the end of the century.<sup>32</sup> Model projections range from small increases to increases of up to 30% in wet areas, and from small decreases to decreases of up to 60% in dry areas.<sup>33 34</sup>
- Generally, windward sides of the major islands will become cloudier and wetter. The dry leeward sides will generally have fewer clouds and less rainfall.<sup>35</sup>

- Stream flow in Hawai'i has declined over approximately the past century, consistent with observed decreases in rainfall.<sup>36</sup> This indicates declining groundwater levels.
- More frequent tropical cyclones are projected for the waters near Hawai'i. This is not necessarily because there will be more storms forming in the east Pacific; rather, it is projected that storms will follow new tracks that bring them into the region of Hawai'i more often.<sup>37</sup>

### **ENSO**

- Frequency of intense El Niño events is projected to double in the 21<sup>st</sup> century, with the likelihood of extreme events occurring roughly once every decade.<sup>38</sup>
- Models project a near doubling in the frequency of future extreme La Niña events, from one in every 23 years to one in every 13 years. Approximately 75% of the increase occurs in years following extreme El Niño events, thus projecting more frequent swings between opposite extremes from one year to the next.<sup>39</sup>
- Strong El Niño years in Hawai'i bring more hot days, intense rains, windless days, active hurricane seasons, and spikes in sea surface temperature.<sup>40</sup>

### **Forest Ecosystems**

- Hawai'i is home to 31% of the nation's plants and animals listed as threatened or endangered, and less than half of the landscape on the islands is still dominated by native plants. Studies indicate that endemic and endangered birds and plants are highly vulnerable to climate change and are already showing shifting habitats.<sup>41</sup>
- Even under moderate warming, 10 of 21 existing native forest bird species are projected to lose over 50% of their range by 2100. Of those, three may lose their entire ranges and three others are projected to lose more than 90% of their ranges making them of high concern for extinction.<sup>42</sup>
- Warming air temperatures are bringing mosquito-borne diseases to previously safe upland forests, driving several native bird species toward extinction.<sup>43</sup>

### **Ocean Warming, Acidification, and Reefs**

- Globally averaged sea surface temperature (SST) increased by 1.8°F (1.0°C) over the past 100 years. Half of this rise has occurred since the 1990s. North Central Pacific averaged SST trends follow the globally averaged trend. Over the last 5 years almost the entire tropical Pacific, in particular areas along the equator, have seen temperatures warmer than the 30 year average.<sup>44</sup>
- Nearly 30 years of oceanic pH measurements, based on data collected from Station ALOHA, Hawai'i, show a roughly 8.7% increase in ocean acidity over this time.<sup>45</sup>
- Increasing ocean acidification reduces the ability of marine organisms to build shells and other hard

structures. This adversely impacts coral reefs and threatens marine ecosystems more broadly.<sup>46</sup>

- In Hawai'i, extended periods of coral bleaching did not first occur until 2014 and 2015 as part of the 2014–17 global scale bleaching event that was the longest ever recorded.<sup>47</sup>
- Ocean warming and acidification are projected to cause annual coral bleaching in some areas, like the central equatorial Pacific Ocean, as early as 2030 and almost all reefs by 2050.<sup>48</sup> This will not only devastate local coral reef ecosystems but will also have profound impacts on ocean ecosystems in general. Ultimately it will threaten the human communities and economies that depend on a healthy ocean.<sup>49</sup>

### **Sea Level Change**

- The mean sea level trend at the Honolulu tide station is 0.055 in (1.41 mm) per year with a 95% confidence interval of  $\pm 0.008$  in (0.21 mm) per year based on monthly mean sea level data, 1905 to 2015. This is equivalent to a change of 0.46 ft (14.0 cm) over the past century.<sup>50</sup>
- The frequency of high tide flooding in Honolulu since the 1960's, has increased from 6 days per year to 11 per year.<sup>51</sup>
- With 3.2 ft (0.98 m) of sea level rise, 25,800 acres experience chronic flooding, erosion, and/or high wave impacts. One third of this land is designated for urban use. Impacts include 38 mi (61 km) of major roads, and more than \$19 billion in assets.<sup>52</sup>
- Due to global gravitational effects, estimates of future sea level rise in Hawai'i and other Pacific islands are about 20%–30% higher than the global mean.<sup>53</sup>
- Over 70% of beaches in Hawai'i are in a state of chronic erosion.<sup>54</sup> This is likely related to long term sea level rise as well as coastal hardening.<sup>55 56</sup>
- Coastal hardening of chronically eroding beaches caused the combined loss of 9% (13.4 mi, 21.5 km) of the length of sandy beaches on Kaua'i, O'ahu, and Maui.<sup>57</sup>

### **Indigenous Communities**

- Indigenous populations will be disproportionately impacted by climate change due to their strong ties to place and greater reliance on natural resources for sustenance.<sup>58</sup>
- About 550 Hawaiian cultural sites are exposed to chronic flooding with a sea level rise of 3.2 ft (0.98 m).<sup>59</sup>
- Sea level rise impacts on traditional and customary practices (including fishpond maintenance, cultivation of salt, and gathering from the nearshore fisheries) have been observed.<sup>60</sup>
- Because of flooding and sea level rise, indigenous practitioners have had limited access to the land where salt is traditionally cultivated and harvested since 2014.

Detachment from traditional lands has a negative effect on the spiritual and mental health of the people.<sup>61</sup>

- Ocean warming and acidification, sea level rise and coastal erosion, drought, flooding, pollution, increased storminess, and over-development are negatively affecting tourism, fisheries, and forested ecosystems. This directly impacts the livelihood and security of Pacific communities. For example, across all Pacific Island countries and territories, industrial tuna fisheries account for half of all exports, 25,000 jobs, and 11% of economic production.<sup>62</sup> In Hawai'i, between 2011 and 2015, an annual average of 37,386 Native Hawaiians worked in tourism-intensive industries; based on the 2013 U.S. census, this number represents 12.5% of the Native Hawaiian population residing in Hawai'i.<sup>63</sup>
- In Hawai'i, climate change impacts, such as reduced streamflow, sea level rise, saltwater intrusion, episodes of intense rainfall, and long periods of drought, threaten the ongoing cultivation of taro and other traditional crops.<sup>64</sup>

### **GLOBAL ECOSYSTEMS**

- Climate change impacts have been documented across every ecosystem on Earth, including shifts in species ranges, shrinking body size, changes in predator-prey relationships, new spawning and seasonal patterns, and modifications in the population and age structure of marine and terrestrial species.<sup>65</sup>
- In 2017 over 15,000 scientists published a "Warning to Humanity".<sup>66</sup> They said humans have pushed Earth's ecosystems to their breaking point and are well on the way to ruining the planet.
- Human activities have increased the acidity of oceans; increased the acidity of freshwater bodies and soils because of acid rain; increased acidity of freshwater streams and groundwater due to drainage from mines; and increased acidity of soils due to added nitrogen to crop lands.<sup>67</sup>
- Researchers have labeled ecosystem impacts "biological annihilation," and identify that a "sixth major mass extinction" is underway as a result of dwindling population sizes and range shrinkages among vertebrates.<sup>68</sup>
- Humans are causing the climate to change 170 times faster than natural forces.<sup>69</sup>
- Tree lines are shifting poleward and to higher elevations.<sup>70</sup>
- One-third of burnt forests experience no tree regeneration at all.<sup>71</sup>
- Species are migrating poleward and to higher elevations.<sup>72</sup>
- Spring is coming sooner to some plant species in the Arctic while other species are delaying their emergence amid warm winters. The changes are associated with diminishing sea ice.<sup>73</sup>

- Spring is coming earlier.<sup>74</sup>
- The tropics have expanded.<sup>75</sup>
- Warmer winters with less snow have resulted in a longer lag time between spring events and a more protracted vernal window (the transition from winter to spring).<sup>76</sup>
- Plants are leafing out and blooming earlier each year.<sup>77</sup>
- Climate-related local extinctions have already occurred in hundreds of species, including 47% of 976 species surveyed.<sup>78</sup>
- Plant and animal extinctions, already widespread, are projected to increase from twofold to fivefold in the coming decades.<sup>79</sup>

## FOOD AND HUMAN HEALTH

- Harvests of staple cereal crops, such as rice and maize, could decline by 20 to 40% as a function of increased surface temperatures in tropical and subtropical regions by 2100.<sup>80</sup>
- One billion people are classified as food insecure.<sup>81</sup>
- Rising CO<sub>2</sub> decreases the nutrient and protein content of wheat, leading to a 15% decline in yield by mid-century.<sup>82</sup>
- Higher levels of CO<sub>2</sub> are lowering amounts of protein, iron, zinc, and B vitamins in rice with potential consequences for a global population of approximately 600 million.<sup>83</sup>
- By 2050, climate change will lead to per-person reductions of 3% in global food availability, 4% in fruit and vegetable consumption, and 0.7% in red meat consumption. These changes will be associated with 529,000 climate-related deaths worldwide.<sup>84</sup>
- Without changes to policy and improvements to technology, food productivity in 2050 could look like it did in 1980 because, at present rates of innovation, new technologies won't be able to keep up with the damage caused by the climate change in major growing regions.<sup>85</sup>
- Certain groups of Americans—including children, elders, the sick and the poor—are most likely to be harmed by climate change.<sup>86</sup>
- Climate change is harming human health now. These harms include heat-related illness, worsening chronic illnesses, injuries and deaths from dangerous weather events, infectious diseases spread by mosquitoes and ticks, illnesses from contaminated food and water, and mental health problems.<sup>87</sup>
- Warming of Earth's surface is changing life on a global scale.<sup>88</sup>
- Extreme weather events are more frequent.<sup>92</sup>
- Half a degree Celsius of global warming has been enough to increase heat waves and heavy rains in many regions of the planet.<sup>93</sup>
- Storm tracks are shifting poleward.<sup>94</sup>
- The number of weather disasters is up 14% since 1995-2004, and has doubled since 1985-1994.<sup>95</sup>
- There has been a global increase in the frequency and intensity of extremely hot three day periods.<sup>96</sup>
- The number of unusually cold days and nights has decreased, and the number of unusually warm days and nights has increased.<sup>97</sup>
- Extreme heat waves are projected to cover double the amount of global land by 2020 and quadruple by 2040, regardless of future emissions trends.<sup>98</sup>
- New records continue to be set for warm temperature extremes. For instance, in the U.S. during February, 2017 there were 3,146 record highs set compared to only 27 record lows, a ratio of 116 to 1.<sup>99</sup>
- Nine of the ten deadliest heat waves have occurred since 2000 causing 128,885 deaths around the world.<sup>100</sup>
- Nearly one third of the world's population is now exposed to climatic conditions that produce deadly heat waves.<sup>101</sup>
- Extreme weather is increasing.<sup>102</sup>
- If global temperatures rise 3.6°F (2°C), the combined effect of heat and humidity will turn summer into one long heat wave. Temperature will exceed 104°F (40°C) every year in many parts of Asia, Australia, Northern Africa, South and North America.<sup>103</sup>
- If global temperatures rise 7.2°F (4°C), a new "super-heatwave" will appear with temperatures peaking at above 131°F making large parts of the planet unlivable including densely populated areas such as the US east coast, coastal China, large parts of India and South America.<sup>104</sup>

## EXTREME WEATHER

- The global percentage of land area in drought has increased about 10%.<sup>89</sup>
- The global occurrence of extreme rainfall has increased 12%.<sup>90</sup>
- Heavy downpours are more intense and frequent.<sup>91</sup>

## GLACIERS, SEA ICE, PERMAFROST

- The world's major ice systems including Antarctica and Greenland,<sup>105</sup> and the mountain glaciers<sup>106</sup> of the world are all in a state of decline.<sup>107, 108, 109</sup>
- Over the past three million years, when global temperatures increased 1.8 to 5.4°F (1 to 3°C), melting polar ice sheets caused global sea levels to rise at least 20 ft (6 m) above present levels.<sup>110</sup>
- Under high emission pathways, a sea level rise exceeding 8 ft (2.4 m) by 2100 is physically possible.<sup>111</sup>
- The West Antarctic ice sheet is in "unstoppable" retreat.<sup>112</sup>
- Atmospheric warming that exceeds 2.7 to 3.6°F (1.5 to 2°C) above present (ca. 2015) will trigger a centennial-to-millennial-scale response of the Antarctic ice sheet that produces an unstoppable contribution to sea-level rise.<sup>113</sup> Substantial Antarctic ice loss can be prevented only by limiting greenhouse gas emissions to RCP2.6

levels. Higher-emissions scenarios lead to ice loss from Antarctica that will raise sea level by 1.9 to 9.8 ft (0.6 to 3 m) by the year 2300.<sup>114</sup>

- If emissions continue unabated, Antarctica has the potential to contribute more than 3.28 ft (1 m) of sea-level rise by 2100 and more than 49.2 ft (15 m) by 2500. In this case, atmospheric warming will soon become the dominant driver of ice loss, but prolonged ocean warming will delay its recovery for thousands of years.<sup>115</sup>
- The Greenland ice sheet is more sensitive to long-term climate change than previously thought. Studies<sup>116</sup> estimate that the warming threshold leading to an essentially ice-free state is in the range of 1.4 to 5.8°F (0.8 to 3.2°C), with a best estimate of 2.9°F (1.6°C) above preindustrial levels. The Arctic is on track to double this amount of warming before mid-century.<sup>117</sup>
- Cloud cover over Greenland is decreasing at 0.9 +/-3% per year. Each 1% of decrease drives an additional 27 +/-13 billion tons of ice melt each year.<sup>118</sup>
- Further melting of mountain glaciers cannot be prevented in the current century - even if all emissions were stopped now.<sup>119</sup> Around 36% of the ice still stored in mountain glaciers today will melt even without further emissions of greenhouse gases. That means: more than one-third of the glacier ice that still exists today in mountain glaciers can no longer be saved even with the most ambitious measures.
- Alpine glaciers have shrunk to their lowest levels in 120 years and are wasting two times faster than they did in the period 1901-1950, three times faster than they did in 1851-1900, and four times faster than they did 1800-1850.<sup>120</sup>
- Arctic sea ice is shrinking (13% per decade) as a result of global warming.<sup>121</sup>
- Winter Arctic sea ice was the lowest on record in 2017.<sup>122</sup>
- In the Arctic, average surface air temperature for the year ending September 2016 was the highest since 1900, and new monthly record highs were recorded for January, February, October, and November 2016.<sup>123</sup>
- Rapid warming in the Arctic is causing the jet stream to slow down and become unstable.<sup>124</sup>
- Regions of Earth where water is frozen for at least one month each year are shrinking with impacts on related ecosystems.<sup>125</sup>
- Extreme warm events in winter are much more prevalent than cold events.<sup>126</sup>
- Global snow cover is shrinking.<sup>127</sup>
- The southern boundary of Northern Hemisphere permafrost is retreating poleward.<sup>128</sup>
- Large parts of permafrost in northwest Canada are slumping and disintegrating into running water. Similar large-scale landscape changes are evident across the Arctic including in Alaska, Siberia, and Scandinavia.<sup>129</sup>

- In North America, spring snow cover extent in the Arctic is the lowest in the satellite record, which started in 1967.<sup>130</sup>

## OCEANS

- The Atlantic Meridional Overturning Circulation has decreased 20%. The North Atlantic has the coldest water in 100 yrs of observations.<sup>131</sup>
- Global sea surface temperature is rising.<sup>132</sup>
- The oceans are warming rapidly.<sup>133</sup>
- Sea level is rising and the rate of rise has accelerated.<sup>134</sup>
- Today global mean sea level is rising three times faster than it was in the 20<sup>th</sup> Century.<sup>135</sup>
- Between 1993 and 2014, the rate of global mean sea level rise increased 50% with the contribution from melting of the Greenland Ice Sheet rising from 5% in 1993 to 25% in 2014.<sup>136</sup>
- With existing greenhouse gas emissions, we are committed to future sea level of at least 4.3 to 6.2 ft (1.3 to 1.9 m) higher than today and are adding about 0.32 m/decade to the total: ten times the rate of observed contemporary sea-level rise.<sup>137</sup>
- Over 90% of the heat trapped by greenhouse gases since the 1970's has been absorbed by the oceans and today the oceans absorb heat at twice the rate they did in the 1990's.<sup>138</sup>
- Excess heat in the oceans has reached deeper waters,<sup>139</sup> and deep ocean temperature is rising.<sup>140</sup>
- Sea surface temperatures have increased in areas of tropical cyclone genesis suggesting a connection with strengthened storminess.<sup>141</sup>
- Oxygen levels in the ocean have declined by 2% over the past five decades because of global warming, probably causing habitat loss for many fish and invertebrate species.<sup>142</sup>
- Marine ecosystems can take thousands, rather than hundreds, of years to recover from climate-related upheavals.<sup>143</sup>
- Marine ecosystems are under extreme stress.<sup>144</sup>
- The world's richest areas for marine biodiversity are also those areas mostly affected by both climate change and industrial fishing.<sup>145</sup>
- The number of coral reefs impacted by bleaching has tripled over the period 1985-2012.<sup>146</sup>
- By 2050 over 98% of coral reefs will be afflicted by bleaching-level thermal stress each year.<sup>147</sup>
- Scientists have concluded that when seas are hot enough for long enough nothing can protect coral reefs. The only hope for securing a future for coral reefs is urgent and rapid action to reduce global warming.<sup>148</sup>
- Average pH of ocean water fell from 8.21 to 8.10, a 30% increase in acidity. Ocean water is more acidic from dissolved CO<sub>2</sub>, which is negatively affecting marine organisms.<sup>149</sup>

- Dissolved oxygen in the oceans is declining because of warmer water.<sup>150</sup>
- Production of oxygen by photosynthetic marine algae is threatened at higher temperatures.<sup>151</sup>

The likely (66%) range of global temperature increase this century will be a median 5.8°F (3.2°C).<sup>152 153</sup> If greenhouse gas concentrations were stabilized at their current level, existing concentrations would commit the world to at least an additional 1.1°F (0.6°C) of warming this century.<sup>154</sup> Beyond the next few decades, the magnitude of climate change depends on emissions of greenhouse gases and aerosols and the sensitivity of the climate system. Projected changes range from 4.7° to 8.6°F (2.6° to 4.8°C) under a higher scenario, to 0.5° to 1.3°F (0.3° to 1.7°C) under the lowest scenario.<sup>155</sup> CO<sub>2</sub> concentration has now passed 400 ppm, a level not seen since 3 million years ago, when global temperature and sea level were significantly higher than today. Testing revealed most climate models underestimate the effects of anthropogenic greenhouse gases.<sup>156</sup> Models that do the best job of simulating observed climate change predict some of the worst-case scenarios for the future. If countries stay on a high-emissions trajectory, there is a 93% chance the planet will warm more than 4°C by the end of the century. Previous studies placed those odds at 62%.

What will this >5.4°F (3°C) world look like?

- Heat waves drive a global scale refugee crisis, as low-latitude continental lands lose habitability<sup>157</sup>;
- Drought<sup>158</sup>, wildfires<sup>159</sup>, water scarcity<sup>160</sup>, crop failure<sup>161</sup> and other threats to critical resources leading to increased human conflict<sup>162</sup>;
- Multi-meter sea level rise continuing over many centuries<sup>163</sup>;
- Extreme weather disasters<sup>164</sup>, massive floods<sup>165</sup>, great tropical cyclones<sup>166</sup>, mega-drought<sup>167</sup>, and torrential rainfall<sup>168</sup> will be widespread.

Ironically, with the ongoing global revolution in clean power, all this suffering and dystopia will be taking place in a world of solar panels, wind mills, electric cars, and cleaner air.

## ENERGY OUTLOOK

### Global

The U.S. Energy Information Administration projects the following global energy patterns to the year 2040.<sup>169</sup>

- Strong, long-term economic growth drives an increasing demand for energy;
- World energy consumption grows by 28%;
- China and India alone account for over half of this increase;
- Fossil fuels maintain a market share of 77% through 2040, even though renewable energy experiences explosive growth;
- World energy-related carbon dioxide emissions rise 15% by 2040.

To hold global temperature below an increase of 3.6°F (2°C) per the 2015 Paris Agreement, it is necessary to decrease carbon emissions by 50% per decade.<sup>170</sup> Clearly, the projections by the EIA move in the opposite direction and present a massive challenge to humanity.

### Hawai'i

What is Hawaii's contribution to greenhouse gas emissions?

- In 2007, Hawaii's total greenhouse gas emissions were 24 million metric tons of CO<sub>2</sub> equivalent;<sup>171</sup>
- Total CO<sub>2</sub> emissions have slightly declined in the last decade, largely due to gains in the electricity sector.<sup>172</sup>

O'ahu had 20.8% of net sales of electricity from sources deemed renewable in 2017, the law requires 100% by 2045.<sup>173</sup>


- Fossil fuel use for transportation continues to increase;<sup>174</sup>
- Hawaii's CO<sub>2</sub> emissions are 20% lower than the national average;<sup>175</sup>
- However, U.S. CO<sub>2</sub> emissions per capita are over three times the world average and Hawaii's are approximately 12 times larger than other Pacific Islands;<sup>176</sup>
- Passed in 2018, Act 15 establishes a *Greenhouse Gas Sequestration Task Force* and sets a 2023 deadline for crafting a plan to meet a zero emissions target by 2045.
- Also passed in 2018, Act 16 directs the state Office of Planning to work with the task force to create a *carbon offset program*.




- <sup>1</sup> Hansen, J., et al. (2016) Ice melt, sea level rise and superstorms: evidence from paleoclimate data, climate modeling, and modern observations that 2°C global warming could be dangerous. *Atmos. Chem. Phys.*, 16, 3761-3812, <https://doi.org/10.5194/acp-16-3761-2016>.
- <sup>2</sup> Keener, V.W., et al. (in review, 2018) Chapter 27, Hawai'i and Pacific Islands, Fourth National Climate Assessment, U.S. Global Change Research Program.
- <sup>3</sup> Islam, N.S. and Winkel, J. (2017) Climate Change and Social Inequality, UN-DESA Work Paper 152, [http://www.un.org/esa/desa/papers/2017/wp152\\_2017.pdf](http://www.un.org/esa/desa/papers/2017/wp152_2017.pdf).
- <sup>4</sup> Ullah, H., et al. (2018) Climate change could drive marine food web collapse through altered trophic flows and cyanobacterial proliferation. *PLOS Biology*; 16 (1): e2003446 DOI: 10.1371/journal.pbio.2003446.
- <sup>5</sup> Barnosky, A.D., et al. (2011) Has the Earth's sixth mass extinction already arrived? *Nature*, 471, 51-57 (03 March).
- <sup>6</sup> Barnosky, A.D., et al. (2012) Approaching a state shift in Earth's biosphere, *Nature*, 486, 7 June, 52-58.
- <sup>7</sup> NOAA, Earth System Research Laboratory, Global Monitoring Division, Frequently Asked questions: [https://www.esrl.noaa.gov/gmd/ccgg/faq\\_cat-3.html](https://www.esrl.noaa.gov/gmd/ccgg/faq_cat-3.html).
- <sup>8</sup> Tripathi, A.K., et al. (2009) Coupling of CO<sub>2</sub> and ice sheet stability over major climate transitions of the last 20 million years, *Science*, 326(5958), 1394-1397.
- <sup>9</sup> Zeebe, R.E., et al. (2016) Anthropogenic carbon release rate unprecedented during the past 66 million years, *Nature Geoscience*, doi: 10.1038/ngeo2681.
- <sup>10</sup> Haustein, K. et al. (2017) A global warming index. *Nature Scientific Reports*, doi:10.1038/s41598-017-14828-5.
- <sup>11</sup> Rafferty, A.E., et al. (2017) Less than 2°C warming by 2100 unlikely, *Nature Climate Change*, 7, 637-641, DOI:10.1038/nclimate3352.
- <sup>12</sup> Hoffman, J.S., et al. (2017) Regional and global SST's during the last interglaciation, *Science*, 355(6322), 276-279, doi: 10.1126/science.aai8464.
- <sup>13</sup> Kopp, R.E., et al. (2009) Probabilistic assessment of sea level during the last interglacial stage, *Nature*, 462, 863-867, doi: 10.1038/nature08686.
- <sup>14</sup> Dutton, A., et al. (2015) Sea-level rise due to polar ice-sheet mass loss during past warm periods, *Science*, v. 349, Is. 6244, DOI: 10.1126/science.aaa4019.
- <sup>15</sup> Willett, K., et al. (2007) Attribution of observed surface humidity changes to human influence, *Nature*, 449, 710-712, doi: 10.1038/nature06207.
- <sup>16</sup> Durack, P., et al. (2012) Ocean salinities reveal strong global water cycle intensification during 1950 to 2000, *Science*, 336(6080), 455-458.
- <sup>17</sup> NOAA National Climatic Data Center, "State of the Climate: Global Analysis for May 2011," published online June 2011, <http://www.ncdc.noaa.gov/sotc/global/>
- <sup>18</sup> Giambelluca, T.W., et al. (2008) Secular Temperature Changes in Hawai'i, *Geophysical Research Letters*, 35:L12702.
- <sup>19</sup> McKenzie, M.M. (2016) Regional temperature trends in Hawai'i: A century of change, 1916-2015 (MS thesis). Dept. of Geog., University of Hawai'i at Mānoa.
- <sup>20</sup> See for example: University of Hawai'i Sea Grant College Program (2014) Climate Change Impacts in Hawai'i - A summary of climate change and its impacts to Hawai'i's ecosystems and communities. UNHI-SEAGRANT-TT-12-04.
- <sup>21</sup> New York Times weather chart: [https://www.nytimes.com/interactive/2016/02/19/us/2015-year-in-weather-temperature-precipitation.html#honolulu\\_hi](https://www.nytimes.com/interactive/2016/02/19/us/2015-year-in-weather-temperature-precipitation.html#honolulu_hi).
- <sup>22</sup> <http://www.hawaiinewsnow.com/story/26551141/hawaiian-electric-asks-oahu-customers-to-conserve-power-tonight>
- <sup>23</sup> Zhang, C., et al. (2016) Dynamical downscaling of the climate for the Hawaiian Islands. Part II: Projection for the late twenty-first century, *Journal of Climate* 29: 8333-8354. doi:10.1175/JCLI-D-16-0038.1.
- <sup>24</sup> Timm, O.E. (2017) Future Warming Rates over the Hawaiian Islands Based on Elevation-Dependent Scaling Factors. *Int. J. Clim.*, doi:10.1002/joc.5065.
- <sup>25</sup> Marra, J.J., and Kruk, M.C. (2017) State of Environmental Conditions in Hawai'i and the U.S. Affiliated Pacific Islands under a Changing Climate: [https://coralreefwatch.noaa.gov/satellite/publications/state\\_of\\_the\\_environment\\_2017\\_hawaii-usapi\\_noaa-nesdis-ncei\\_oct2017.pdf](https://coralreefwatch.noaa.gov/satellite/publications/state_of_the_environment_2017_hawaii-usapi_noaa-nesdis-ncei_oct2017.pdf).
- <sup>26</sup> Marra and Kruk (2017)
- <sup>27</sup> Storlazzi, C.D., et al. (2015). Future wave and wind proj. for US and US-API. USGS OFR No. 2015-1001, <http://dx.doi.org/10.3133/ofr20151001>.
- <sup>28</sup> Frazier, A.G. and Giambelluca, T.W. (2017) Spatial trend analysis of HI rainfall from 1920 to 2012. *Int. J. Climatol.*, 37: 2522-2531, DOI: 10.1002/joc.4862.
- <sup>29</sup> Frazier and Giambelluca (2017)
- <sup>30</sup> Kruk, M. C., et al. (2015), On the state of the knowledge of rainfall extremes in the western and northern Pacific basin, *Int. J. Climatol.*, 35(3), 321-336.
- <sup>31</sup> Kruk et al. (2015)
- <sup>32</sup> PIRCA (2016) Expert Consensus on Downscaled Climate Projections for the Main Hawaiian Islands. PIRCA Information Sheet, HI. <http://bit.ly/2yoY0II>.
- <sup>33</sup> Zhang et al (2016)
- <sup>34</sup> Timm, O.E., et al. (2015), Stat. downscaling of rainfall changes in HI, based on CMIP5 model proj., *JGR Atmos*, 120(1), 92-112, doi:10.1002/2014JD022059.
- <sup>35</sup> Zhang et al (2016)
- <sup>36</sup> Bassiouni, M., and D.S. Oki. 2013. Trends and shifts in stream flow in Hawai'i, 1913-2008. *Hydrological Processes* 27(10):1484-1500.
- <sup>37</sup> Murakami, H., et al. (2013) Projected increase in tropical cyclones near Hawai'i. *Nature Climate Change*, v. 3, August, pp. 749-754.
- <sup>38</sup> Cai, W., et al. (2015) Inc. frequency of extreme El Niño events due to greenhouse warming. *Nature Climate Change* 4, 111-116, doi:10.1038/nclimate2100.
- <sup>39</sup> Cai, W., et al. (2015) Inc. freq. of extreme La Niña events induced by greenhouse warming, *Nature Climate Change*, 5, 132-137, doi: 10.1038/nclimate2492.
- <sup>40</sup> Keener et al. (in review, 2018)
- <sup>41</sup> Jacobi, J.D., et al. (2017) Baseline land cover. In Selmants, P.C., et al., eds., USGS, <http://pubs.er.usgs.gov/publication/pp1834>.
- <sup>42</sup> Fortini, L., et al. (2015) Large-scale range collapse of HI forest birds under CC and the need 21st century conservation options. *PLoS ONE*, 10.
- <sup>43</sup> Paxton, E.H., et al. (2016) Collapsing avian community on a Hawaiian island. *Science Advances*, 2, e1600029.
- <sup>44</sup> Marra and Kruk (2017)
- <sup>45</sup> Marra and Kruk (2017)
- <sup>46</sup> Marra and Kruk (2017)
- <sup>47</sup> Marra and Kruk (2017)
- <sup>48</sup> Van Hooijdonk, R., et al. (2014) Opposite latitudinal gradients in projected ocean acidification and bleaching impacts on coral reefs. *Global Change Biology*, 20.
- <sup>49</sup> Marra and Kruk (2017)
- <sup>50</sup> Marra and Kruk (2017)
- <sup>51</sup> Marra and Kruk (2017)
- <sup>52</sup> Hawai'i Sea Level Rise Vulnerability and Adaptation Report (2017) Tetra Tech, Inc. and the State of Hawai'i DLNR, OCCL, DLNR Contract No: 64064.
- <sup>53</sup> Marra and Kruk (2017)
- <sup>54</sup> Fletcher, C.H., et al. (2012) National Assessment of Shoreline Change: Historical shoreline change in the Hawaiian Islands. USGS OFR 2011-1051, 55p.
- <sup>55</sup> Romine, B.M., et al. (2013) Are beach erosion rates and sea-level rise related in Hawai'i? *Global and Planetary Change*, 108: 149-157.
- <sup>56</sup> Romine, B.M. and Fletcher, C.H. (2012) Armoring on eroding coasts leads to beach narrowing and loss on O'ahu, HI. DOI 10.1007/978-94-007-4123-2\_10.
- <sup>57</sup> Fletcher et al. (2012)
- <sup>58</sup> Climate Home News (2014) 5 ways CC harms indigenous people. <http://www.climatechangenews.com/2014/07/28/five-ways-climate-change-harms-indigenous-people/>.
- <sup>59</sup> Hawai'i Climate Change Mitigation and Adaptation Commission (2017).
- <sup>60</sup> Sproat, D. K. (2016) An Indigenous People's Right to Environmental Self-Determination: Native Hawaiians and the Struggle Against Climate Change Devastation. *Stanford Environmental Law Journal*, 35.
- <sup>61</sup> Akutagawa, M., et al. (2016) Health Impact Assessment of the Proposed Mo'omomi Community-Based Subsistence Fishing Area (Report). The Kohala Center.
- <sup>62</sup> Gillett, R., et al. (2001) Tuna: a key economic resource in the Pacific Islands. Pacific studies series. Manila, Philippines: Asian Development Bank.

- <sup>63</sup> Keener et al. (in review, 2018)
- <sup>64</sup> Sproat (2016)
- <sup>65</sup> Scheffers, B.R., et al. (2016) The broad footprint of climate change from genes to biomes to people. *Science*, Nov., DOI: 10.1126/science.aaf7671.
- <sup>66</sup> Ripple, W.J., et al. (2017) World Scientists' Warning to Humanity: A Second Notice. *BioScience*.
- <sup>67</sup> Rice, K. and Herman, J. (2012) Acidification of Earth: an assessment across mechanisms and scales, *Applied Geochemistry*, 27(1), 1-14.
- <sup>68</sup> Ceballos, G., et al. (2017) Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines, *PNAS*, 114 (30) E6089-E6096; doi:10.1073/pnas.1704949114
- <sup>69</sup> Gaffney, O., and Steffen, W. (2017) The Anthropocene equation, *The Anthropocene Review*, <http://dx.doi.org/10.1177%2F2053019616688022>
- <sup>70</sup> Beck, P.S.A., et al. (2011) Changes in forest productivity across Alaska consistent with biome shift, *Ecology Letters*, doi: 10.1111/j.1461-0248.2011.01598.x
- <sup>71</sup> Stevens-Rumann, C.S., et al. (2018) Evidence for declining forest resilience to wildfires under climate change, *Ecology Letters*, 21:243-252.
- <sup>72</sup> Loarie, S.R., et al. (2009) The velocity of climate change, *Nature*, 462, 1052-1055.
- <sup>73</sup> Post, E., et al. (2016) Highly individualistic rates of plant phenological advance associated with arctic sea ice dynamics, *Biology Letters*, 12(12), 20160332.
- <sup>74</sup> The U.S. Geological Survey hails an early spring and ties it to climate change: <http://www.chron.com/news/houston-weather/article/The-U-S-Geological-Survey-hails-an-early-spring-10958042.php>. Kahru, M., et al. (2010) Are phytoplankton blooms occurring earlier in the Arctic? *Global Change Biology*, doi: 10.1111/j.1365-2486.2010.02312.x.
- <sup>75</sup> Seidel, D.J., et al. (2008) Widening of the tropical belt in a changing climate, *Nature Geoscience*, 1, 21-24, doi: 10.1038/ngeo.2007.38.
- <sup>76</sup> A. R. Contosta, et al. (2017) A longer vernal window: the role of winter coldness and snowpack in driving spring transitions and lags. *Global Change Biology*, 23 (4): 1610 DOI: 10.1111/gcb.13517.
- <sup>77</sup> Wolkovich, E., et al. (2012) Warming experiments underpredict plant phenological responses to climate change, *Nature*, 485(7399), 494-497.
- <sup>78</sup> Wiens, J.J. (2016) Climate-related local extinctions are already widespread among plant and animal species, *PLOS Biology*, 14(12), e2001104.
- <sup>79</sup> Wiens, J.J. (2016)
- <sup>80</sup> Battisti, D.S. and Naylor, R.L. (2009) Historical warnings of future food insecurity with unprecedented seasonal heat. *Science* 323, 240–244,
- <sup>81</sup> Barrett, C.B. (2010) Measuring food insecurity. *Science* 327, 825–828.
- <sup>82</sup> Myers, S.S., et al. (2014) Increasing CO<sub>2</sub> threatens human nutrition, *Nature*, 510, 139-142, doi: 10.1038/nature13179. Feng, Z., et al. (2015) Constraints to nitrogen acquisition of terrestrial plants under elevated CO<sub>2</sub>, *Global Change Biology*, 21(8), 3152-3168, doi: 10.1111/gcb.12938.
- <sup>83</sup> Zhu, C., et al. (2018) Carbon dioxide (CO<sub>2</sub>) levels this century will alter the protein, micronutrients, and vitamin content of rice grains with potential health consequences for the poorest rice-dependent countries, *Science Advances*, 23 May, v. 4, no. 5, DOI:10.1126/sciadv.aag1012.
- <sup>84</sup> Springmann, M., et al. (2016) Global and regional health effects of future food production under climate change: a modeling study, *The Lancet*.
- <sup>85</sup> Liang, X.Z., et al. (2017) Determining climate effects on US total agricultural productivity, *PNAS*, [www.pnas.org/cgi/doi/10.1073/pnas.1615922114](http://www.pnas.org/cgi/doi/10.1073/pnas.1615922114)
- <sup>86</sup> Medical Alert! Climate change is Harming Our Health, report by the Medical Society Consortium on Climate and Health, 24p. [https://medsocietiesforclimatehealth.org/wp-content/uploads/2017/03/medical\\_alert.pdf](https://medsocietiesforclimatehealth.org/wp-content/uploads/2017/03/medical_alert.pdf).
- <sup>87</sup> U.S Global Change Research Program (2016) The Impacts of Climate Change on Human Health in the United States: A Scientific Assessment. Crimmins, A., et al. GCRP, Washington, DC, 312 pp. <http://dx.doi.org/10.7930/J0R49NQX>.
- <sup>88</sup> Rosenzweig, C., et al. (2008) Attributing Physical and Biological Impacts to Anthropogenic Climate Change, *Nature* 453, no. 7193: 353–357.
- <sup>89</sup> Dai, A. (2011) Characteristics and trends in various forms of the Palmer drought severity index during 1900–2008, *Journal of Geophysical Research* 116.
- <sup>90</sup> Lehmann, J., et al. (2015) Increased record-breaking precipitation events under global warming. *Climatic Change*, doi: 10.1007/s10584-015-1434-y
- <sup>91</sup> See NOAA, <https://www.climate.gov/news-features/featured-images/heavy-downpours-more-intense-frequent-warmer-world>.
- <sup>92</sup> Medvigy, D. and Beaulieu, C. (2011) Trends in daily solar radiation and precipitation coefficients of variation since 1984, *Journal of Climate*, 25(4), 1330-1339.
- <sup>93</sup> Schleussner, C-F, et al. (2017) In the observational record half a degree matters, *Nature Climate Change*. DOI: 10.1038/nclimate3320.
- <sup>94</sup> Bender, F. A-M, et al. (2012) Changes in extratropical storm track cloudiness 1983–2008: Observational support for a poleward shift, *Climate Dynamics* 38.
- <sup>95</sup> Centre for Research on the Epidemiology of Disasters, UN International Strategy for Disaster Reduction, <http://reliefweb.int/report/world/human-cost-weather-related-disasters-1995-2015>.
- <sup>96</sup> Wehner, M., et al. (2018) Early 21<sup>st</sup> Century anthropogenic changes in extremely hot days as simulated by the C20C+ detection and attribution multi-model ensemble, *Weather and Climate Extremes*, v. 20, June, p. 1-8, <https://doi.org/10.1016/j.wace.2018.03.001>.
- <sup>97</sup> Easterling, D.R., et al. (2016) Detection and attribution of climate extremes in the observed record, *Weather and Climate Extremes*, 11, March, p. 17-27.
- <sup>98</sup> Coumou, D. and Robinson, A. (2013) Historic and future increase in the global land area affected by monthly heat extremes, *Environmental Research Letters*, 8(3), 034018, doi: 10.1088/1748-9326/8/3/034018.
- <sup>99</sup> See Climate Central.org, <http://www.climatecentral.org/news/record-high-temperature-february-21186>.
- <sup>100</sup> Vaidyanathan, G. (2015) Killer heat grows hotter around the world, *Scientific American*, August 6.
- <sup>101</sup> Mora, C. et al. (2017) Global risk of deadly heat. *Nature Climate Change*; DOI: 10.1038/NCLIMATE332.
- <sup>102</sup> S. Rahmstorf and D. Coumou (2011) Increase in Extreme Events in a Warming World. *PNAS* 108, no. 44: 17905–17909. Francis, J.A., and Vavrus, S.J. (2012) Evidence linking Arctic amplification to extreme weather in mid-latitudes. *GRL*, 39, L06801. Stott, P. (2016) How climate change affects extreme weather events. *Science*, 352, 1517–1518, doi:10.1126/science.aaf7271.
- <sup>103</sup> Russo, S., et al. (2017) Humid heat waves at different warming levels. *Scientific Reports*; 7 (1) DOI: 10.1038/s41598-017-07536-7.
- <sup>104</sup> Russo, S., et al. (2017)
- <sup>105</sup> Data from NASA's GRACE satellites show that the land ice sheets in both Antarctica and Greenland have been losing mass since 2002. Both ice sheets have seen an acceleration of ice mass loss since 2009: <https://climate.nasa.gov>.
- <sup>106</sup> Marzeion, B., et al. (2018) Limited influence of climate change mitigation on short-term glacier mass loss, *Nature Climate Change*.
- <sup>107</sup> Radić, V. and Hock, R. (2011) Regionally differentiated contribution of mountain glaciers and ice caps to future sea-level rise, *Nature Geoscience*, 4, 91-94.
- <sup>108</sup> Zemp, M., et al. (2015) Historically unprecedented global glacier decline in the early 21st century. *Journal of Glaciology*; 61 (228): 745.
- <sup>109</sup> Rignot, E., et al. (2011) Acceleration of the contribution of the Greenland and Antarctic ice sheets to sea level rise, *Geophysical Research Letters*, 38, L05503.
- <sup>110</sup> Dutton, A., et al. (2015) Sea-level rise due to polar ice-sheet mass loss during past warm periods, *Science*, 10 Jul., v. 349, ls. 6244.
- <sup>111</sup> USGCRP (2017)
- <sup>112</sup> Joughlin, I., et al. (2014) Marine ice sheet collapse potentially underway for the Thwaites Glacier Basin, West Antarctica, *Science*, May 12. Rignot, E., et al. (2014) Widespread, rapid grounding line retreat of Pine Island, Thwaites, Smith and Kohler glaciers, West Antarctica from 1992 to 2011, *GRL*.
- <sup>113</sup> Colledge, N.R., et al. (2015) The multi-millennial Antarctic commitment to future sea-level rise, *Nature*, 526 (7573): 421 DOI: 10.1038/nature15706.
- <sup>114</sup> Colledge et al. (2015)
- <sup>115</sup> DeConto, R.M. and Pollard, D. (2016) Contribution of Antarctica to past and future sea-level rise, *Nature*, 531 (7596):591–597.
- <sup>116</sup> Robinson, A., et al. (2012) Multistability and Critical Thresholds of the Greenland Ice Sheet, *Nature Climate Change*, 2, 429–432.
- <sup>117</sup> Smith, S.J., et al. (2015) Near-term acceleration in the rate of temperature change, *Nature Climate Change*, March 9, DOI: 10.1038/nclimate2552.


- 118 Hofer, S., et al. (2017) Decreasing cloud cover drives the recent mass loss on the Greenland ice sheet. *Science Advances*, 28 June, v. 3, no. 6, e1700584.
- 119 Marzeion et al. (2018)
- 120 Zemp et al. (2015)
- 121 Arctic Report Card: <http://www.arctic.noaa.gov/Report-Card/Report-Card-2016>. Serreze, M., et al. (2007) Perspectives on the Arctic's shrinking sea-ice cover, *Science* 315, 1533-1536.
- 122 See <http://nsidc.org/arcticseaicenews/>.
- 123 See <http://www.arctic.noaa.gov/Report-Card/Report-Card-2016>.
- 124 Francis, J., and Skific, N. (2015) Evidence linking rapid Arctic warming to mid-latitude weather patterns, *Phil. Trans. R. Soc. A*, 373, 20140170.
- 125 Fountain, A., et al. (2012) The disappearing cryosphere: Impacts and ecosystem responses to rapid cryosphere loss, *BioScience* 62(4), 405-415.
- 126 Guirguis, K., et al. (2011) Recent warm and cold daily winter temperature extremes in the northern hemisphere, *Geophysical Research Letters*, 38, L17701.
- 127 Déry, S. J. and Brown, R.D. (2007) Recent NH snow cover extent trends and implications for the snow albedo feedback, *GRL*, 34, L22504
- 128 Thibault, S. and Payette, S. (2009) Recent permafrost degradation in bogs of the James Bay area, Northern Quebec, Canada, *Permafrost and Periglacial Processes*, 20(4), 383, doi: 10.1002/ppp.660.
- 129 Kokelj, S.V., et al. (2017) Climate-driven thaw of permafrost preserved glacial landscapes, northwestern Canada, *Geology*, Feb. doi: 10.1130/G38626.1
- 130 See Arctic Report Card, <http://www.arctic.noaa.gov/Report-Card/Report-Card-2016>
- 131 Rahmstorf, S., et al. (2015) Exceptional twentieth-century slowdown in Atlantic Ocean overturning circulation. *Nature Climate Change*.
- 132 Levitus, S., et al. (2008) Global ocean heat content in light of recently revealed instrumentation problems, *GRL*, 36, L07608, doi: 10.1029/2008GL037155
- 133 Wang, G., et al. (2017) Consensuses and discrepancies of basin-scale ocean heat content changes in different ocean analyses, *Climate Dynamics*.
- 134 Nerem, R.S., et al. (2018) Climate-change-driven accelerated sea-level rise detected in the altimeter era. *PNAS*, DOI: 10.1073/pnas.1717312115.
- 135 Dangendorf, S., et al. (2017) Reassessment of 20<sup>th</sup> Century global mean sea level rise, *PNAS*, doi: 10.1073/pnas.1616007114
- 136 Chen, X., et al. (2017) The increasing rate of global mean sea-level rise during 1993–2014, *Nature Climate Change*. DOI: 10.1038/nclimate3325
- 137 Strauss, B.H. (2015) Rapid accumulation of committed sea level rise from global warming, *PNAS*, 110(34), 13699-13700.
- 138 Cheng L., et al (2015) Global upper ocean heat content estimation: recent progress and the remaining challenges. *Atmospheric and Oceanic Science Letters*, 8, DOI:10.3878/AOSL20150031. Glecker, P.J., et al. (2016) Industrial era global ocean heat uptake doubles in recent decades. *Nature Climate Change*.
- 139 Cheng, L., et al. (2017) Improved estimates of ocean heat content from 1960 to 2015, *Science Advances* 10 Mar., v. 3, no. 3, e1601545.
- 140 Song, Y.T. and Colberg, F. (2011) Deep ocean warming assessed from altimeters, GRACE in situ measurements, and a non-Boussinesq ocean general circulation model, *JGR* 116, C02020. Volkov, D.L., et al. (2017) Decade-long deep-ocean warming detected in the subtropical South Pacific, *GRL*.
- 141 Defforge, C.L. and Merlis, T.M. (2017) Observed warming trend in sea surface temperature at tropical cyclone genesis, *GRL*.
- 142 Schmidtko, S., et al. (2017) Decline in global oceanic oxygen content during the past five decades, *Nature*, 542, 335-339, 16 February.
- 143 S.E. Moffitt, et al. (2015) Response of seafloor ecosystems to abrupt global climate change. *PNAS*, 2015 DOI: 10.1073/pnas.1417130112.
- 144 McCauley, D.J., et al. (2015) Marine defaunation: Animal loss in the global ocean, *Science*, 347(6219), 16, Jan. Henson, S.A., et al. (2017) Rapid emergence of climate change in environmental drivers of marine ecosystems, *Nature Communications*, 8, 14682, doi: 10.1038/ncomms14682.
- 145 Ramirez, F., et al. (2017) Climate impacts on global hot spots of marine biodiversity. *Science Advances*; 3 (2): e1601198 DOI: 10.1126/sciadv.1601198.
- 146 Heron, S.F., et al. (2016) Warming trends and bleaching stress of the worlds coral reefs 1985-2012, *Scientific Reports*, 6, 38402, doi: 10.1038/srep38402.
- 147 Heron, S.F., et al. (2016)
- 148 Hughes, T.P., et al. (2017) Global warming and recurrent mass bleaching of corals. *Nature*; 543 (7645): 373 DOI: 10.1038/nature21707.
- 149 Barton, A., et al. (2012) The Pacific oyster, *Crossostrea gigas*, shows negative correlation to naturally elevated carbon dioxide levels: Implications for near-term ocean acidification effects, *Limnology and Oceanography* 57(3), 698-710, doi: 10.4319/lo.2012.57.3.0698.
- 150 L. Stramma, et al. (2011) Expansion of Oxygen Minimum Zones May Reduce Available Habitat for Tropical Pelagic Fishes, *Nature Climate Change* 2: 33–37.
- 151 Sekerci, Y. and Petrovskii (2015) Mathematical modeling of Plankton-Oxygen dynamics under the climate change. *Bulletin of Mathematical Biology*.
- 152 Raftery et al. (2017)
- 153 Tollefson, J. (2018) Can the world kick its fossil fuel addiction fast enough? *Nature*, News Feature, 25 April.
- 154 Hayhoe, K., et al. (2017) Climate models, scenarios, and projections. In: CSSR: 4NCA, v. I [Wuebbles, D.J., et al. (eds.)]. USGCRP, Wash., DC.
- 155 Hayhoe et al. (2017)
- 156 Brown, P.T., Caldeira, K. (2017) Greater future global warming inferred from Earth's recent energy budget, *Nature* 552, 45-50, DOI:10.1038/nature24672.
- 157 Russo, S., et al. (2017); Mora, C., et al. (2017)
- 158 Dai, A. (2013) Increasing drought under global warming in observations and models, *Nature Climate Change*, 3, p. 52-58, doi:10.1038/nclimate1633.
- 159 Jolly, W.M., et al. (2015) Climate-induced variations in global wildfire danger from 1979 to 2013, *Nature Communications* 6, DOI:10.1038/ncomms8537.
- 160 Richey, A. S., et al. (2015) Quantifying renewable groundwater stress with GRACE, *Water Resour. Res.*, 51, 5217–5238, doi:10.1002/2015WR017349.
- 161 Zhao, C., et al. (2017) Temperature increase reduces global yields of major crops in four independent estimates *PNAS* 114 (35) 9326-9331.
- 162 Raleigh, C., et al. (2014) Extreme temperatures and violence. *Nature Climate Change*, 4, 76–77.
- 163 Dutton et al. (2015) See also Golledge, N.R., et al. (2015) The multi-millennial Antarctic commitment to future sea-level rise: *Nature*, 2015; 526 (7573): 421.
- 164 Centre for Res. on the Epidemiology of Disasters, UN Intl. Strat. for Disaster Red.: <http://reliefweb.int/report/world/human-cost-weather-related-disasters-1995-2015>
- 165 Arnell, N.W., and Gosling, S.N. (2016) The impacts of climate change on river flood risk at the global scale, *Climatic Change*, 134, 3, pp 387-401.
- 166 Kossin, J.P., et al. (2017) Extreme storms. In: CSSR: NCA4, v I [Wuebbles, D.J., et al., eds.] USGCRP, Wash, DC, pp. 257-276, doi: 10.7930/J07S7KXX.
- 167 Ault, T.R., et al. (2016) Relative impacts of mitigation, temperature, and precipitation on 21<sup>st</sup> Century megadrought risk in the American Southwest. *Science Advances*, Oct 5, v. 2, no. 10, DOI:10.1126/sciadv.1600873.
- 168 Lehmann, J., et al. (2015) Increased record-breaking precipitation events under global warming. *Climatic Change*, doi: 10.1007/s10584-015-1434-y
- 169 EIA (2017) International Energy Outlook. U.S. Energy Information Administration, [https://www.eia.gov/outlooks/ieo/pdf/0484\(2017\).pdf](https://www.eia.gov/outlooks/ieo/pdf/0484(2017).pdf)
- 170 Rockström, J. et al. (2017) A roadmap for rapid decarbonization. *Science*, 355 (6331): 1269. Gasser, T., et al. (2015) Negative emissions physically needed to keep global warming below 2°C, *Nature Communications* 6, DOI: 10.1038/ncomms8958. Carbon Brief (2016) Explainer: 10 ways negative emissions could slow climate change: <https://www.carbonbrief.org/explainer-10-ways-negative-emissions-could-slow-climate-change>
- 171 Hawaii Greenhouse Gas Inventory. 1990 and 2007. <https://energy.hawaii.gov/wp-content/uploads/2011/10/ghg-inventory-20081.pdf>
- 172 State Carbon Dioxide Emissions Data. <https://www.eia.gov/environment/emissions/state/>
- 173 Renewable Portfolio Standards Law Examination, Docket No. 2007-0008. <https://puc.hawaii.gov/wp-content/uploads/2018/02/RPS-HECO-2017.pdf>
- 174 State Energy Data System (SEDS) 1960-2015. <https://www.eia.gov/state/seds/seds-data-complete.php?sid=HI#CompleteDataFile>
- 175 State Carbon Dioxide Emissions Data. <https://www.eia.gov/environment/emissions/state/>
- U.S. Department of Commerce, Census Bureau. <https://www.census.gov/programs-surveys/popest/data/tables.html>
- 176 CO2 emissions (metric tons per capita). [https://data.worldbank.org/indicator/EN.ATM.CO2E.PC?year\\_high\\_desc=false](https://data.worldbank.org/indicator/EN.ATM.CO2E.PC?year_high_desc=false)



**Figure 2.** Climate models indicate that the world can stop global warming at 2°C if greenhouse gas emissions are reduced by 50% per decade immediately. Delaying this reduction leads to warming of more than 3°C – a dangerous level. Source: Levin, K., and Rich, D. (2017) *Turning Points: Trends in Countries' Reaching Peak Greenhouse Gas Emissions over Time*. Working Paper. Washington, DC: World Resources Institute. Available online at <http://www.wri.org/publication/turning-points>.



**Figure 3.** Although consumption of nonfossil fuels is expected to grow faster than fossil fuels, fossil fuels still account for 77% of energy use in 2040. Source: U.S. Energy Information Administration, *International Energy Outlook 2017, Executive Summary*: [https://www.eia.gov/outlooks/ieo/exec\\_summ.php](https://www.eia.gov/outlooks/ieo/exec_summ.php)



**Figure 4.** Although some large emitters, including China, the EU and India have either reduced—or slowed—their GHG emissions growth rate, currently implemented policies are expected to result in a further growth of global GHG emissions by about 9–13% between 2020–2030. Source: Climate Action Tracker: <https://climateactiontracker.org/publications/improvement-warming-outlook-india-and-china-move-ahead-paris-agreement-gap-still-looms-large/>